

ORIGINAL ARTICLE

Intravenous Vitamin C in Adults with Sepsis in the Intensive Care Unit

F. Lamontagne, M.-H. Masse, J. Menard, S. Sprague, R. Pinto, D.K. Heyland, D.J. Cook, M.-C. Battista, A.G. Day, G.H. Guyatt, S. Kanji, R. Parke, S.P. McGuinness, B.-K. Tirupakuzhi Vijayaraghavan, D. Annane, D. Cohen, Y.M. Arabi, B. Bolduc, N. Marinoff, B. Rochweg, T. Millen, M.O. Meade, L. Hand, I. Watpool, R. Porteous, P.J. Young, F. D'Aragon, E.P. Belley-Cote, E. Carbonneau, F. Clarke, D.M. Maslove, M. Hunt, M. Chassé, M. Lebrasseur, F. Lauzier, S. Mehta, H. Quiroz-Martinez, O.G. Rewa, E. Charbonney, A.J.E. Seely, D.J. Kutsogiannis, R. LeBlanc, A. Mekontso-Dessap, T.S. Mele, A.F. Turgeon, G. Wood, S.S. Kohli, J. Shahin, P. Twardowski, and N.K.J. Adhikari, for the LOVIT Investigators and the Canadian Critical Care Trials Group*

ABSTRACT

BACKGROUND

Studies that have evaluated the use of intravenous vitamin C in adults with sepsis who were receiving vasopressor therapy in the intensive care unit (ICU) have shown mixed results with respect to the risk of death and organ dysfunction.

METHODS

In this randomized, placebo-controlled trial, we assigned adults who had been in the ICU for no longer than 24 hours, who had proven or suspected infection as the main diagnosis, and who were receiving a vasopressor to receive an infusion of either vitamin C (at a dose of 50 mg per kilogram of body weight) or matched placebo administered every 6 hours for up to 96 hours. The primary outcome was a composite of death or persistent organ dysfunction (defined by the use of vasopressors, invasive mechanical ventilation, or new renal-replacement therapy) on day 28.

RESULTS

A total of 872 patients underwent randomization (435 to the vitamin C group and 437 to the control group). The primary outcome occurred in 191 of 429 patients (44.5%) in the vitamin C group and in 167 of 434 patients (38.5%) in the control group (risk ratio, 1.21; 95% confidence interval [CI], 1.04 to 1.40; $P=0.01$). At 28 days, death had occurred in 152 of 429 patients (35.4%) in the vitamin C group and in 137 of 434 patients (31.6%) in the placebo group (risk ratio, 1.17; 95% CI, 0.98 to 1.40) and persistent organ dysfunction in 39 of 429 patients (9.1%) and 30 of 434 patients (6.9%), respectively (risk ratio, 1.30; 95% CI, 0.83 to 2.05). Findings were similar in the two groups regarding organ-dysfunction scores, biomarkers, 6-month survival, health-related quality of life, stage 3 acute kidney injury, and hypoglycemic episodes. In the vitamin C group, one patient had a severe hypoglycemic episode and another had a serious anaphylaxis event.

CONCLUSIONS

In adults with sepsis receiving vasopressor therapy in the ICU, those who received intravenous vitamin C had a higher risk of death or persistent organ dysfunction at 28 days than those who received placebo. (Funded by the Lotte and John Hecht Memorial Foundation; LOVIT ClinicalTrials.gov number, NCT03680274.)

The authors' full names, academic degrees, and affiliations are listed in the Appendix. Dr. Lamontagne can be contacted at francois.lamontagne@usherbrooke.ca or at the Department of Medicine, Université de Sherbrooke, 3001 12th Ave. N., Sherbrooke, Quebec J1H 5N4, Canada. Dr. Adhikari can be contacted at neil.adhikari@utoronto.ca or at the Department of Critical Care Medicine, Sunnybrook Health Sciences Centre, 2075 Bayview Ave., Room D1.08, Toronto M4N 3M5, Canada.

*A complete list of the LOVIT investigators is provided in the Supplementary Appendix, available at NEJM.org.

Drs. Lamontagne and Adhikari contributed equally to this article.

This article was published on June 15, 2022, at NEJM.org.

N Engl J Med 2022;386:2387-98.

DOI: 10.1056/NEJMoa2200644

Copyright © 2022 Massachusetts Medical Society.

 A Quick Take
is available at
NEJM.org

SEPSIS IS DEFINED AS LIFE-THREATENING organ dysfunction caused by a dysregulated host response to infection.¹ This disorder causes or contributes to between a third and a half of deaths in hospitals² and is responsible for as many as 11 million deaths worldwide each year.³ Treatment includes antimicrobial therapy, source control, and organ support.

In sepsis, the antioxidant effects of vitamin C therapy⁴ may mitigate tissue injury induced by oxidative stress.⁵ Vitamin C cannot be synthesized by humans, and levels are low in many critically ill patients, which has increased the plausibility of benefit with supplementation.⁶ After a single-center study spurred interest in the use of intravenous vitamin C, administered with hydrocortisone and thiamine,⁷ subsequent randomized, controlled trials evaluating this combination treatment did not show benefits.^{8,9} In contrast, in a randomized, controlled trial, patients with sepsis and acute lung injury who received a higher dose of vitamin C (50 mg per kilogram of body weight every 6 hours) had a lower 28-day risk of death than those who received placebo.¹⁰ However, recent meta-analyses suggest that the overall evidence supporting the use of vitamin C therapy in patients with sepsis is of low certainty.^{8,11}

In the phase 3, multicenter, randomized, controlled Lessening Organ Dysfunction with Vitamin C (LOVIT) trial, we tested the hypothesis that a high dose of vitamin C would reduce the risk of death or persistent organ dysfunction at 28 days in adults with sepsis who were receiving vasopressor therapy in the intensive care unit (ICU).

METHODS

TRIAL DESIGN

In this international trial, we enrolled patients in 35 adult medical–surgical ICUs in **Canada, France, and New Zealand**. The protocol (available with the full text of this article at NEJM.org) has been described previously^{12,13} and was approved by the ethics committee at each participating trial site.

The trial was funded by the Lotte and John Hecht Memorial Foundation. Nova Biomedical Canada provided glucometers, testing strips, and control solutions (StatStrip Express) to trial

sites that requested them. Without input from the funder, the authors were responsible for the design, planning, and coordination of the trial and for the analysis of the data; all the authors made the decision to submit the manuscript for publication. Site investigators, research personnel, or trained delegates assessed the eligibility of potential patients, and research personnel collected the data. Informed consent was provided by the patients or their legal representatives; after approval by local authorities, consent could be obtained by telephone or patients could be enrolled with deferred consent, followed by informed consent as soon as reasonably possible.

The corresponding authors wrote the first draft of the manuscript. All the authors vouch for the accuracy and completeness of the data and for the fidelity of the trial to the protocol.

PATIENTS

Eligible patients were **adults (≥18 years of age) who had been in the ICU for no longer than 24 hours, who had proven or suspected infection as the main diagnosis, and who were receiving a vasopressor**. Exclusion criteria included contraindications to vitamin C therapy, receipt of open-label vitamin C, or expected death or withdrawal of life-sustaining therapy within 48 hours. Details regarding the inclusion and exclusion criteria are provided in the Supplementary Appendix, available at NEJM.org.

RANDOMIZATION AND TREATMENT

We randomly assigned patients in a 1:1 ratio to receive either **vitamin C or placebo**, stratified according to site by means of a centralized Web-based system using permuted blocks of variable, undisclosed size. Patients, clinicians, and trial personnel and statisticians were unaware of trial-group assignments.

Patients in the intervention group received intravenous vitamin C in a bolus dose of 50 mg per kilogram mixed in a 50-ml solution of either dextrose 5% in water or normal saline. Doses were administered over 30 to 60 minutes every 6 hours for 96 hours (i.e., 200 mg per kilogram per day, with a maximum of 16 doses) as long as patients remained in the ICU. In the control group, patients received a matching placebo infusion (dextrose 5% in water or normal saline). At each site, pharmacists who were not involved

in the patients' clinical care prepared the respective infusions in an unblinded manner.

All other aspects of care, including the administration of glucocorticoids (including agents with mineralocorticoid effects) and thiamine, were performed at the discretion of the treating teams. Local research staff members collected data on inpatient outcomes, and the central management team conducted telephone interviews with patients or their representatives 6 months after randomization. If patients had been discharged from the hospital before day 28, ascertainment of the primary outcome was completed at the time of the 6-month follow-up interview.

TRIAL OUTCOMES

The primary outcome was a composite of death or persistent organ dysfunction (defined as receipt of vasopressors, invasive mechanical ventilation, or new renal-replacement therapy)¹⁴ on trial day 28. Secondary outcomes were the number of days without organ dysfunction in the ICU up to day 28; mortality at 28 days and 6 months; quality of life at 6 months; organ failure at days 2, 3, 4, 7, 10, 14, and 28; and biomarkers of global tissue dysoxia (lactate), inflammation (interleukin-1 β and tumor necrosis factor α), and endothelial injury (thrombomodulin and angiotensin-2) at days 3 and 7. Quality of life was assessed with the use of the European Quality of Life-5 Dimension 5-Level (EQ-5D-5L) questionnaire,¹⁵ which evaluates mobility, personal care, usual activities, pain or discomfort, and anxiety or depression and categorizes each of these dimensions into five levels that range from no problems to extreme problems. Organ failure was measured by means of the score on the Sequential Organ Failure Assessment (SOFA),¹⁶ which grades the function of six organ systems on the basis of blood pressure and vasopressor requirements, oxygenation, platelet count, serum creatinine and bilirubin levels, and the score on the Glasgow Coma Scale. We evaluated the patients' disease severity on the Acute Physiology and Chronic Health Evaluation (APACHE) II, with scores that range from 0 to 71, with higher scores indicating an increased risk of death.

On the basis of potential adverse effects reportedly associated with vitamin C therapy,¹⁷⁻²⁰ we recorded the incidence of stage 3 acute kidney injury,^{17,18} acute hemolysis,¹⁹ and hypoglycemia²⁰ as safety outcomes.

All unexpected serious adverse events (i.e., those neither prespecified nor included as outcomes) that were considered by the investigator to be at least possibly related to a trial procedure were reported to the trial coordinating center within 24 hours and were subsequently investigated and reported to the data and safety monitoring board and to Health Canada, the governmental department responsible for Canadian health policy.

STATISTICAL ANALYSIS

On the basis of the results of a clinical trial involving a similar population,²¹ we anticipated that the risk of death at 28 days or persistent organ dysfunction in the control group would be approximately 50%. Thus, the enrollment of 385 patients per group would provide the trial with 80% power to detect an absolute between-group difference of 10 percentage points in the risk of this outcome with a two-sided type I error rate (α) of 0.05. To account for withdrawal of consent and loss to follow-up, we planned to enroll 400 patients per group. On April 23, 2020, the steering committee notified the data and safety monitoring board and Health Canada that patients with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection who fulfilled the eligibility criteria would be offered participation in the trial. We inflated the sample size to ensure that the trial included the originally intended number of patients without SARS-CoV-2 infection.

The primary analysis was performed in the intention-to-treat population to assess the superiority of vitamin C over placebo, according to the assigned trial group. We estimated the risk ratio and 95% confidence interval for the primary outcome in a generalized linear mixed model with binomial distribution and a log-link function, with trial site considered as a random effect.²² In a secondary analysis of the primary outcome, we adjusted for prespecified baseline characteristics (age, sex, APACHE II score,²³ baseline receipt of glucocorticoids, and time from ICU admission to randomization) using generalized estimating equations.

In other prespecified secondary analyses, we compared mortality at 28 days in unadjusted and adjusted models by applying the same variables that were used in analyzing the primary out-

come. We compared 6-month survival using a Cox model and compared the number of days free of organ dysfunction in the ICU up to day 28 using a cumulative distribution function, with death at 28 days coded as minus one. In addition, we compared SOFA scores during the first 7 days using a linear mixed model that included time, group interaction, and biomarkers according to constrained longitudinal data analysis.²⁴ For SOFA scores after day 7, we evaluated between-group differences in means and imputed scores for patients who had died or had been discharged before day 7. For safety outcomes, we reported the number of each prespecified safety outcome (after adjudication in the case of hemolysis) and unexpected serious adverse events in each trial group. We report treatment-effect estimates as risk ratios or differences in means or medians as appropriate, with no adjustment for multiple comparisons.

Analyses excluded one patient for whom outcome data were missing after withdrawal of consent for follow-up. We conducted best case–worst case unadjusted sensitivity analyses of death as a component of the primary outcome and as a secondary outcome.

The data and safety monitoring board reviewed the results of two planned interim analyses (after 248 and 525 patients had completed follow-up for the primary outcome) and recommended continuation of the trial. Details regarding all statistical analyses are provided in the protocol^{12,13} and the Supplementary Appendix. At the time of this report, all the trial investigators remained unaware of the results of the interim analyses.

We prespecified the subgroup analyses of the primary outcome according to age (<65 years or ≥65 years), sex, frailty (according to a score on the Clinical Frailty Scale²⁵ of 1 to 4 or ≥5), severity of illness (the quartile of predicted risk of death on the basis of the baseline APACHE II score), presence of septic shock (defined as the use of a vasopressor infusion to maintain a mean arterial pressure of ≥65 mm Hg and the presence of a lactate level of ≥2 mmol per liter¹ vs. the use of vasopressor infusion alone), and the quartile of the baseline vitamin C level (as measured by liquid chromatography–tandem mass spectrometry). We hypothesized that vitamin C would be more beneficial in elderly pa-

Figure 1 (facing page). Enrollment and Randomization.

Immediately after randomization but before receiving any dose of either vitamin C or placebo, 8 patients were found to be ineligible and were removed from the trial after blinded adjudication by two steering-committee members. Of these patients, 1 underwent randomization under deferred consent but regained sufficient capacity to decline consent; 5 patients had surrogate decision makers who initially gave and then withdrew consent, 1 patient was found to be allergic to vitamin C, and 1 patient was enrolled in error in a different study using the same randomization system. In addition, 1 patient withdrew consent after receiving the trial infusion. This patient remained part of the trial population according to the intention-to-treat principle. G6PD denotes glucose-6-phosphate dehydrogenase, ICU intensive care unit, and IV intravenous.

tients, in those with greater frailty and illness severity at baseline, in those who met strict criteria for septic shock, and in those with a lower baseline vitamin C level. After modifying the protocol to permit the enrollment of patients with SARS-CoV-2, we added a subgroup analysis comparing the effect of vitamin C in patients with and without SARS-CoV-2 that was based on the hypothesis of no difference in treatment effect. In addition, we assessed the credibility of apparent subgroup effects.²⁶

RESULTS

PATIENTS

From November 14, 2018, to July 19, 2021, we enrolled 872 patients; of these patients, 8 underwent randomization in error and 1 withdrew consent, which left 863 patients in the primary analysis population (429 in the vitamin C group and 434 in the placebo group) (Fig. 1).

The characteristics of the patients at baseline were similar in the two groups (Table 1 and Table S1 in the Supplementary Appendix). Overall, 96.7% of the patients received at least 90% of the scheduled doses of vitamin C or placebo (Table S2). The patients' median stay was 6 days (interquartile range [IQR], 3 to 12) in the ICU and 16 days (IQR, 8 to 32) in the hospital. The use and duration of cointerventions and life-sustaining therapies during the course of the ICU stay were similar in the two groups (Tables S3 and S4). The groups had similar urine output and fluid balance during the first 7 days in the ICU (Table S5).




Table 1. Characteristics of the Patients at Baseline.*

Characteristic	Vitamin C (N = 429)	Placebo (N = 433)†
Age — yr	65.0±14.0	65.2±13.8
Female sex — no. (%)	151 (35.2)	173 (40.0)
Admission type — no. (%)‡		
Medical	350 (81.6)	369 (85.2)
Emergency surgery	69 (16.1)	59 (13.6)
Elective surgery	10 (2.3)	5 (1.2)
APACHE II score§	24.2±7.4	24.1±7.9
SOFA score¶	10.2±3.4	10.1±3.7
Score on Clinical Frailty Scale	3.8±1.4	3.9±1.4
1 to 4 — no. (%)	312 (72.7)	308 (71.3)
≥5 — no. (%)	117 (27.3)	124 (28.7)
Primary site of infection — no. (%)**		
Pulmonary	145 (33.8)	159 (36.7)
Gastrointestinal or intra-abdominal	133 (31.0)	112 (25.9)
Blood	55 (12.8)	59 (13.6)
Skin or soft tissue	55 (12.8)	62 (14.3)
Urinary	49 (11.4)	55 (12.7)
Central nervous system	2 (0.5)	4 (0.9)
Other	30 (7.0)	27 (6.2)
SARS-CoV-2 positive — no. (%)††	37 (8.6)	26 (6.0)
Lactate — mmol/liter‡‡	3.4±3.2	3.0±2.8
Vitamin C — μmol/liter§§	20.6±70.6	19.1±39.7
Septic shock definition met — no./total no. (%)¶¶	195/327 (59.6)	183/326 (56.1)
Time from ICU admission to randomization — hr	12.9±8.2	12.3±6.7
Treatment — no. (%)		
Glucocorticoid	199 (46.4)	196 (45.4)
Mechanical ventilation	294 (68.5)	283 (65.4)
Renal-replacement therapy	46 (10.7)	42 (9.7)
Vasopressor infusion****	428 (99.8)	433 (100)

* Plus–minus values are means ±SD.

† One additional patient underwent randomization under deferred consent and died before consent was obtained. The research ethics board allowed the inclusion of this patient only in the primary analysis.

‡ Patients with emergency or elective surgical admission came to the intensive care unit (ICU) from the operating room or postanesthetic care unit.

§ Scores on the Acute Physiologic and Chronic Health Evaluation (APACHE) II range from 0 to 71, with higher scores indicating greater disease severity and a higher risk of death.

¶ Scores on the Sequential Organ Failure Assessment (SOFA) were calculated from the worst values corresponding to six organ systems on day 1 (day of randomization). As such, the worst values could precede or follow randomization and the first trial infusion. Scores range from 0 to 24, with higher scores indicating more severe organ dysfunction.

|| Scores on the Clinical Frailty Scale range from 1 to 9, with higher scores indicating greater frailty. One patient in the placebo group had missing data.

** Patients could have more than one site of infection.

†† In these patients, coronavirus disease 2019 was confirmed or suspected at baseline and subsequently confirmed.

‡‡ Data were available for 653 patients (327 in the vitamin C group and 326 in the placebo group).

§§ The normal value for vitamin C in plasma is 22.4 μmol per liter (range, 13.6 to 32.9).¹⁸ Data were available for 646 patients (324 in the vitamin C group and 322 in the placebo group).

¶¶ The definition for septic shock included the requirement for a vasopressor infusion and a lactate level of at least 2 mmol per liter.

||| Data were missing for 1 patient in the placebo group.

**** Vasopressor infusion was discontinued in 1 patient between the time that consent had been obtained and randomization.

Table 2. Primary and Secondary Outcomes.*

Outcome	Vitamin C	Placebo	Treatment Effect (95% CI) [†]
Primary			
Death or persistent organ dysfunction at 28 days — no./total no. (%) [‡]	191/429 (44.5)	167/434 (38.5)	1.21 (1.04 to 1.40)
Death	152/429 (35.4)	137/434 (31.6)	1.17 (0.98 to 1.40)
Persistent organ dysfunction [§]	39/429 (9.1)	30/434 (6.9)	1.30 (0.83 to 2.05)
Vasopressor use	8/429 (1.9)	6/434 (1.4)	1.36 (0.48 to 3.85)
Mechanical ventilation	25/429 (5.8)	19/434 (4.4)	1.31 (0.74 to 2.30)
Renal-replacement therapy	24/429 (5.6)	18/434 (4.1)	1.35 (0.73 to 2.5)
Secondary			
Median no. of days without organ dysfunction in the ICU by day 28 (IQR) [¶]	17 (–1 to 25)	19.5 (–1 to 25)	–2.43 (–7.23 to 2.37)
Death by 6 mo — no./total no. (%)	191/417 (45.8)	185/426 (43.4)	1.14 (0.93 to 1.39)
EQ-5D-5L score at 6 mo ^{¶¶} **			
Score on visual-analogue scale	65.8±20.9	63.8±22.5	2.04 (–1.97 to 6.05)
Median dimension score (IQR)			
Mobility	2 (1–3)	2 (1–3)	–0.19 (–0.43 to 0.04)
Self-care	1 (1–2)	1 (1–2)	–0.07 (–0.29 to 0.15)
Usual activities	2 (2–4)	2 (2–3)	0.02 (–0.23 to 0.28)
Pain or discomfort	2 (1–3)	2 (1–3)	0.00 (–0.19 to 0.18)
Anxiety or depression	1 (1–2)	2 (1–2)	–0.08 (–0.24 to 0.09)
SOFA score ^{¶¶¶}			
Day 1 (N=862)	10.2±3.4	10.1±3.7	0.05 (–0.42 to 0.53)
Day 2 (N=862)	9.9±5.2	9.5±5.1	0.39 (–0.30 to 1.07)
Day 3 (N=861)	9.2±6.0	9.0±6.0	0.23 (–0.57 to 1.03)
Day 4 (N=861)	8.7±6.5	8.7±6.6	–0.03 (–0.90 to 0.85)
Day 7 (N=862)	9.0±7.9	8.3±7.3	0.66 (–0.35 to 1.67)
Day 10 (N=277)	7.5±4.4	7.5±3.9	0.05 (–0.94 to 1.05)
Day 14 (N=191)	7.4±4.2	7.3±4.2	0.07 (–1.12 to 1.26)
Day 28 (N=56)	6.5±3.8	7.9±5.7	–1.42 (–3.98 to 1.14)
Safety outcomes — no./total no. (%)^{‡‡}			
Stage 3 acute kidney injury	162/429 (37.8)	164/433 (37.9) ^{§§}	1.00 (0.85 to 1.19)
Acute hemolysis ^{¶¶}	0	0	NA
Hypoglycemia	26/429 (6.1)	22/433 (5.1) ^{§§}	1.25 (0.73 to 2.14)
Serious adverse events	1/429 (0.2)	0	NA

* Plus–minus values are means ±SD. IQR denotes interquartile range, and NA not applicable.

[†] The treatment effect is a risk ratio unless otherwise indicated. Confidence intervals have not been adjusted for multiplicity, and inferences drawn from the intervals may not be reproducible.

[‡] P=0.01 for this comparison.

[§] The analysis of the individual components of persistent organ dysfunction was added post hoc. Patients could have more than one component.

[¶] The treatment effect in this category is the between-group difference.

^{||} The treatment effect in this category is a hazard ratio.

^{**} Scores on the EQ-5D-5L instrument range from 1 to 5, with higher scores indicating greater impairment or worse function. Scores on the visual-analogue scale in that instrument range from 0 to 100, with higher scores indicating better health status. Data were available for 222 patients in the vitamin C group and for 233 in the placebo group.

^{††} Missing SOFA scores for days 1 to 7 were imputed according to the statistical analysis plan, with the exception of one patient on days 3 and 4 because of withdrawal of consent.

^{‡‡} Additional data regarding safety outcomes are provided in Table S8 in the Supplementary Appendix.

^{§§} One patient who had undergone randomization under deferred consent died before consent was obtained. The research ethics board allowed the inclusion of this patient only in the primary analysis.

^{¶¶} Included in this category are the results of adjudicated events.

^{|||} An anaphylactic reaction in this patient was considered by the investigator to be possibly related to the vitamin C infusion.

SAFETY OUTCOMES AND SUBGROUP ANALYSES

We observed no material between-group differences in the prespecified safety outcomes (Table 2). There were 4 adverse events in the vitamin C group and 1 in the placebo group (Table S8). In the vitamin C group, 1 patient had a serious adverse event of anaphylaxis and another had a severe hypoglycemic episode²⁷ that triggered a protocol amendment requiring modifications in blood-glucose monitoring in many centers. There was no evidence of credible subgroup effects (Fig. 3 and Table S9).




Figure 2. Kaplan–Meier Analysis of Survival at 6 Months.

Shown is the percentage of patients who were alive at the 6-month follow-up (226 patients [54.2%] in the vitamin C group and 241 [56.6%] in the placebo group), which was a secondary outcome in the trial.

DISCUSSION

In this international, randomized, placebo-controlled clinical trial involving adults with sepsis who were receiving vasopressor infusions in the ICU, the composite primary outcome (death or persistent organ dysfunction at trial day 28) occurred more frequently in patients who had received intravenous vitamin C than in those who had received placebo. This was an unexpected finding, and the secondary analyses — which included the evaluation of five biomarkers of tissue dysoxia, inflammation, and endothelial injury measured up to day 7 — did not determine a putative mechanism for harm.

Our findings differ from those of recent meta-analyses of vitamin C monotherapy,^{8,11} which included two smaller randomized trials evaluating the same vitamin C regimen as was used in our trial.^{10,28} The first of these was a dose-finding trial²⁸ involving 24 patients with severe sepsis who were randomly assigned to receive vitamin C at a dose of either 12.5 mg or 50 mg per kilogram or placebo every 6 hours for 4 days.²⁸ The higher-dose vitamin C regimen was associated with a reduction in SOFA scores during a 96-hour period. In the second trial (Vitamin C Infusion for Treatment in Sepsis Induced Acute Lung Injury [CITRIS-ALI]),¹⁰ 167 patients with sepsis and acute respiratory distress syndrome were randomly assigned to receive vitamin C (50 mg per kilogram) or placebo every 6 hours for 4 days. SOFA scores during a 96-hour period were similar in the two groups without accounting for the competing risk of death, but 28-day mortality was significantly lower in the vitamin C group. As in our trial, in the CITRIS-ALI trial, patients had undergone randomization within 24 hours

PRIMARY OUTCOME

At day 28, 191 of 429 patients (44.5%) in the vitamin C group had died or had persistent organ dysfunction, as compared with 167 of 434 patients (38.5%) in the placebo group (risk ratio, 1.21; 95% confidence interval [CI], 1.04 to 1.40; $P=0.01$) (Table 2). In the analysis performed after adjustment for prespecified baseline characteristics, the risk ratio for the primary comparison was 1.15 (95% CI, 0.90 to 1.47) (Table S6). In the best case–worst case sensitivity analyses that accounted for the single patient with unknown status regarding the primary outcome, the results were similar to those in the primary analysis.

SECONDARY OUTCOMES

At 28 days, death had occurred in 152 patients (35.4%) in the vitamin C group and in 137 patients (31.6%) in the placebo group (risk ratio, 1.17; 95% CI, 0.98 to 1.40). The median number of days without organ dysfunction at day 28 was 17 in the vitamin C group and 19.5 in the placebo group (median difference, -2.43 days; 95% CI, -7.23 to 2.37) (Table 2 and Fig. S1). We observed no material differences between groups in SOFA scores (Fig. S2), biomarkers (Table S7), 6-month survival (Fig. 2), or health-related quality of life (Table 2).




Figure 3. Subgroup Analyses.

Shown is a forest plot of the primary outcome in prespecified subgroups. Risk ratios are based on generalized linear mixed models after adjustment for trial site, assigned group, subgroup, and the interaction between subgroup and assigned group. For the subgroup with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), a generalized estimating equation was used because the generalized linear mixed model did not converge. The score on the Clinical Frailty Scale was missing for one patient in the placebo group. Statistical tests for interaction appear in Table S9. Confidence intervals have not been adjusted for multiplicity and inferences drawn from the intervals may not be reproducible. The predicted risk of death was estimated on the basis of the patient's score on the Acute Physiology and Chronic Health Evaluation (APACHE) II, on which higher scores indicate an increased risk of death. Higher scores on the Clinical Frailty Scale indicate more severe frailty. Sepsis-3 denotes the Third International Consensus Definitions for Sepsis and Septic Shock.

after eligibility had been confirmed. However, in our trial, patients did not need to have severe respiratory failure and may have been recruited earlier relative to the onset of sepsis and peak oxidative stress than were the patients in the CITRIS-ALI trial. In our trial, vitamin C was administered within 4 hours after randomization, as compared with 6 hours in the CITRIS-ALI trial.

One plausible explanation for the divergent findings regarding mortality is that large effect estimates from smaller trials may occur by chance.²⁹ Differences in baseline characteristics,

such as the presence of respiratory failure or receipt of vasopressors, may also explain such differences. Such knowledge gaps may be addressed in ongoing trials of vitamin C in patients with SARS-CoV-2 infection (ClinicalTrials.gov numbers NCT02735707 and NCT04401150) and in those with acute respiratory distress syndrome (EudraCT number, 2020-003923-40).

The primary outcome in our trial, a composite of death or persistent organ dysfunction, included death to address the competing-risk issue.¹⁴ Vitamin C had a consistent effect across all elements of the composite outcome and on 6-month

mortality. These factors, in addition to increased statistical efficiency, mitigate the disadvantages of using a composite outcome.^{30,31} Other strengths of our analyses include blinding to limit ascertainment bias, a median enrollment time of approximately 12 hours after ICU admission, high protocol adherence, ascertainment of the renal-replacement therapy component of the primary outcome in patients who were discharged from the hospital before day 28, and assessment of biomarkers and baseline vitamin C levels.

However, the trial also has several limitations. Nine patients who had undergone randomization did not contribute data to the primary analysis; of these patients, eight had not received either vitamin C or placebo and had met the prespecified criteria for exclusion after randomization.³² Only one patient who withdrew consent could not be included in the intention-to-treat analysis. Given the high number of events that were recorded in each group, the effect of these exclusions is probably small. Information regarding specific pathogens and the appropriateness of antimicrobial therapy was not collected; however, systematic differences in post-randomization care are less likely in blinded trials. Information to ascertain the presence of acute respiratory distress syndrome at baseline was not collected, so it remains unclear whether this subgroup had a different response to vitamin C.

Although the patients were representative of those with sepsis being treated in the ICU in many high-income countries, the trial population differs substantially from patients in many low- and middle-income countries, where the incidence of sepsis and the associated case fatality rate are highest (Table S10).³

In adults with sepsis who were receiving vasopressor therapy in the ICU, the receipt of intravenous vitamin C resulted in a higher risk of death or persistent organ dysfunction at 28 days than the receipt of placebo.

Supported by the Lotte and John Hecht Memorial Foundation. Nova Biomedical Canada provided glucometers, testing strips, and control solutions (StatStrip Express) to trial sites that requested them. Vitamin C that was used at trial sites in Canada and New Zealand was purchased from Mylan.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

A data sharing statement provided by the authors is available with the full text of this article at NEJM.org.

We thank the trial patients and their caregivers; the health care providers, research and laboratory staff members, and trial and data coordinators; the staff members at the Unité de Recherche Clinique et Épidémiologique du Centre de Recherche du CHU de Sherbrooke for providing support in the conduct of the trial; Louise Robert Petit for her work with biologic specimens and supplies; Marie-Pier Domingue for her assistance in the preparation of earlier versions of the figures; Anitra Carr for her insights regarding the potential underlying mechanisms of the effects of vitamin C in previous studies; members of the Canadian Critical Care Trials Group for their input on methodologic issues, including Ron Wald for his review of the manuscript; and the members of the data and safety monitoring board (Andreas Laupacis, Lauren Griffith, and Scott Halpern); and the broader Code C community, whose support proved essential to the success of this trial.

APPENDIX

The authors' full names and academic degrees are as follows: François Lamontagne, M.D., Marie-Hélène Masse, M.Sc., Julie Menard, Ph.D., Sheila Sprague, Ph.D., Ruxandra Pinto, Ph.D., Daren K. Heyland, M.D., Deborah J Cook, M.D., Marie-Claude Battista, Ph.D., Andrew G. Day, M.Sc., Gordon H. Guyatt, M.D., Salmaan Kanji, Pharm.D., Rachael Parke, R.N., M.H.Sc., Ph.D., Shay P. McGuinness, M.B., Ch.B., Bharath-Kumar Tirupakuzhi Vijayaraghavan, M.D., Djillali Annane, M.D., Ph.D., Dian Cohen, B.A., Yaseen M. Arabi, M.D., Brigitte Bolduc, M.Sc., Nicole Marinoff, R.N., Bram Rochweg, M.D., Tina Millen, R.R.T., Maureen O. Meade, M.D., Lori Hand, B.Sc., Irene Watpool, R.N., B.Sc.N., Rebecca Porteous, B.N.Sc., Paul J. Young, M.B., Ch.B., Ph.D., Frederick D'Aragnon, M.D., Emilie P. Belley-Cote, M.D., Ph.D., Elaine Carbonneau, R.N., France Clarke, R.R.T., David M. Maslove, M.D., Miranda Hunt, B.A., Michaël Chassé, M.D., Ph.D., Martine Lebrasseur, R.N., François Lauzier, M.D., Sangeeta Mehta, M.D., Hector Quiroz-Martinez, M.D., Oleksa G. Rewa, M.D., Emmanuel Charbonney, M.D., Ph.D., Andrew J.E. Seely, M.D., Ph.D., Demetrios J. Kutsogiannis, M.D., M.H.S., Remi LeBlanc, M.D., Armand Mekontso-Dessap, M.D., Ph.D., Tina S. Mele, M.D., Ph.D., Alexis F. Turgeon, M.D., Gordon Wood, M.D., Sandeep S. Kohli, M.D., Jason Shahin, M.D.C.M., Pawel Twardowski, M.D., Ph.D., and Neill K.J. Adhikari, M.D.C.M.

The authors' affiliations are as follows: the Departments of Medicine (F. Lamontagne, M.-C.B., H.Q.M.) and Anesthesiology (F.D.), Université de Sherbrooke, the Department of Pharmacy (B.B.), Centre de Recherche du CHU de Sherbrooke, Centre Intégré Universitaire de Santé et de Services Sociaux de l'Estrie—Centre Hospitalier Universitaire de Sherbrooke (F. Lamontagne, M.-H.M., J.M., F.D., E. Carbonneau), and Bishop's University (D.C.), Sherbrooke, QC, the Division of Orthopaedic Surgery, Department of Surgery (S.S.), the Department of Medicine (D.J.C., G.H.G., B.R., T.M., M.O.M., E.P.B.-C., S.S.K.), the Department of Health Research Methods, Evidence, and Impact (D.J.C., G.H.G., B.R., M.O.M., L.H., E.P.B.-C., F.C.), and the Population Health Research Institute (E.P.B.-C.), McMaster University, the Department of Critical Care, St. Joseph's Healthcare Hamilton (D.J.C.), the Intensive Care Unit, Juravinski Hospital (T.M.), and Respiratory Therapy, Hamilton Health Sciences (L.H.), Hamilton, ON, the Department of Critical Care Medicine, Sunnybrook Health Sciences Centre (R. Pinto, N.M., N.K.J.A.), the Interdepartmental Division of Critical Care Medicine, University of Toronto (S.M., N.K.J.A.), and Sinai Health System (S.M.), Toronto, the Departments of Critical Care Medicine (D.K.H., D.M.M., M.H.), Public Health Sciences (A.G.D.), and Medicine (D.M.M.), Queen's University, and Kingston Health Sciences Centre (A.G.D., D.M.M.), Kingston, ON, the Department of Pharmacy, Ottawa Hospital (S.K.), Ottawa Hospital Research Institute (S.K., I.W., R. Porteous), and the Department of Surgery, University of Ottawa (A.J.E.S.), Ottawa, ON, the Massawippi Valley Foundation, Ayer's Cliff, QC (D.C.), the

Department of Medicine, Université de Montréal (M.C.), the Centre Hospitalier de l'Université de Montréal (M.C., M.L., E. Charbonney), the Centre de Recherche de l'Hôpital du Sacré-Coeur de Montréal (E. Charbonney), and the Departments of Medicine and Critical Care, McGill University Health Centre (J.S.), Montreal, QC, the Department of Medicine (F. Lauzier) and the Division of Critical Care Medicine, Department of Anesthesiology and Critical Care (F. Lauzier, A.F.T.), Université Laval, the Critical Care Medicine Service (F. Lauzier, A.F.T.), and the Research Center (F. Lauzier, A.F.T.), CHU de Québec—Université Laval, Quebec, QC, the Department of Critical Care Medicine, University of Alberta, Edmonton (O.G.R., D.J.K.), the Division of Intensive Care, Department of Medicine, CHU Dr. Georges L. Dumont, Moncton, NB (R.L.), the Department of Surgery, Schulich School of Medicine and Dentistry, Western University, London, ON (T.S.M.), the Division of Intensive Care Medicine, Island Health Authority, Victoria, BC (G.W.), and the Department of Medicine, Halton Healthcare, Oakville, ON (S.S.K.) — all in Canada; the Cardiothoracic and Vascular Intensive Care Unit, Auckland City Hospital (R. Parke, S.P.M.), and the School of Nursing, University of Auckland (R. Parke), Auckland, the Medical Research Institute of New Zealand, Newtown (R. Parke, S.P.M., P.J.Y.), the Intensive Care Unit, Wellington Hospital, Wellington (P.J.Y.), and the Department of Surgical Sciences, University of Otago, Dunedin (P.T.) — all in New Zealand; the Department of Critical Care Medicine, Apollo Hospitals, Chennai, and George Institute for Global Health, New Delhi — both in India (B.K.T.V.); the Department of Intensive Care Medicine, Raymond-Poincaré Hospital (Assistance Publique—Hôpitaux de Paris [AP-HP]), and Fédération Hospitalo-Universitaire SEPSIS, Garches (D.A.), Institut National de la Santé et de la Recherche Médicale, University Paris-Saclay Campus UVSQ, Versailles (D.A.), and Service de Médecine Intensive Réanimation, AP-HP, Hôpitaux Universitaires Henri-Mondor, Groupe de Recherche Clinique CARMAS, and Institut Mondor de Recherche Biomédicale, Université Paris Est Créteil, Créteil (A.M.-D.) — all in France; the College of Medicine, King Saud Bin Abdulaziz University for Health Sciences, King Abdullah International Medical Research Center, and the Department of Intensive Care, King Abdulaziz Medical City, Ministry of National Guard Health Affairs — all in Riyadh, Saudi Arabia (Y.M.A.); and the Australian and New Zealand Intensive Care Research Centre, Monash University, Melbourne, VIC, Australia (R. Parke, S.P.M., P.J.Y.).

REFERENCES

- Singer M, Deutschman CS, Seymour CW, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 2016;315:801-10.
- Liu V, Escobar GJ, Greene JD, et al. Hospital deaths in patients with sepsis from 2 independent cohorts. *JAMA* 2014;312:90-2.
- Rudd KE, Johnson SC, Agesa KM, et al. Global, regional, and national sepsis incidence and mortality, 1990-2017: analysis for the Global Burden of Disease Study. *Lancet* 2020;395:200-11.
- Frei B, Stocker R, England L, Ames BN. Ascorbate: the most effective antioxidant in human blood plasma. *Adv Exp Med Biol* 1990;264:155-63.
- Amrein K, Oudemans-van Straaten HM, Berger MM. Vitamin therapy in critically ill patients: focus on thiamine, vitamin C, and vitamin D. *Intensive Care Med* 2018;44:1940-4.
- Carr AC, Rosengrave PC, Bayer S, Chambers S, Mehrtens J, Shaw GM. Hypovitaminosis C and vitamin C deficiency in critically ill patients despite recommended enteral and parenteral intakes. *Crit Care* 2017;21:300.
- Marik PE, Khangoora V, Rivera R, Hooper MH, Catravas J. Hydrocortisone, citamin C, and thiamine for the treatment of severe sepsis and septic shock: a retrospective before-after study. *Chest* 2017;151:1229-38.
- Fujii T, Salanti G, Belletti A, et al. Effect of adjunctive vitamin C, glucocorticoids, and vitamin B1 on longer-term mortality in adults with sepsis or septic shock: a systematic review and a component network meta-analysis. *Intensive Care Med* 2022;48:16-24.
- Sevransky JE, Rothman RE, Hager DN, et al. Effect of vitamin C, thiamine, and hydrocortisone on ventilator- and vasopressor-free days in patients with sepsis: the VICTAS randomized clinical trial. *JAMA* 2021;325:742-50.
- Fowler AA III, Truitt JD, Hite RD, et al. Effect of vitamin C infusion on organ failure and biomarkers of inflammation and vascular injury in patients with sepsis and severe acute respiratory failure: the CITRIS-ALI randomized clinical trial. *JAMA* 2019;322:1261-70.
- Patel JJ, Ortiz-Reyes A, Dhaliwal R, et al. IV vitamin C in critically ill patients: a systematic review and meta-analysis. *Crit Care Med* 2022;50(3):e304-e312.
- Masse MH, Ménard J, Sprague S, et al. Lessening Organ dysfunction with Vitamin C (LOVIT): protocol for a randomized controlled trial. *Trials* 2020;21:42.
- Adhikari NK, Pinto R, Day AG, et al. Lessening Organ dysfunction with Vitamin C (LOVIT): statistical analysis plan. *JMIR Res Protoc* 2022 April 14 (Epub ahead of print).
- Heyland DK, Muscedere J, Drover J, Jiang X, Day AG. Persistent organ dysfunction plus death: a novel, composite outcome measure for critical care trials. *Crit Care* 2011;15:R98.
- Herdman M, Gudex C, Lloyd A, et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res* 2011;20:1727-36.
- Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med* 1996;22:707-10.
- Kellum JA, Lameire N, Aspelin P, et al. KDIGO clinical practice guideline for acute kidney injury. *Kidney Int Suppl* 2012;2:Suppl 1:1-38.
- de Groot H-J, Manubulu-Choo W-P, Zandvliet AS, et al. Vitamin C pharmacokinetics in critically ill patients: a randomized trial of four IV regimens. *Chest* 2018;153:1368-77.
- Rees DC, Kelsey H, Richards JD. Acute haemolysis induced by high dose ascorbic acid in glucose-6-phosphate dehydrogenase deficiency. *BMJ* 1993;306:841-2.
- Tang Z, Du X, Louie RF, Kost GJ. Effects of drugs on glucose measurements with handheld glucose meters and a portable glucose analyzer. *Am J Clin Pathol* 2000;113:75-86.
- Heyland D, Muscedere J, Wischmeyer PE, et al. A randomized trial of glutamine and antioxidants in critically ill patients. *N Engl J Med* 2013;368:1489-97.
- Kahan BC. Accounting for centre-effects in multicentre trials with a binary outcome — when, why, and how? *BMC Med Res Methodol* 2014;14:20.
- Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Crit Care Med* 1985;13:818-29.
- Coffman CJ, Edelman D, Woolson RF. To condition or not condition? Analysing 'change' in longitudinal randomised controlled trials. *BMJ Open* 2016;6(12):e013096.
- Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ* 2005;173:489-95.
- Schandelmaier S, Briel M, Varadhan R, et al. Development of the Instrument to assess the Credibility of Effect Modification Analyses (ICEMAN) in randomized controlled trials and meta-analyses. *CMAJ* 2020;192(32):E901-E906.
- Lachance O, Goyer F, Adhikari NKJ, et al. High-dose vitamin-C induced prolonged factitious hyperglycemia in a peri-

toneal dialysis patient: a case report. *J Med Case Rep* 2021;15:297.

28. Fowler AA III, Syed AA, Knowlson S, et al. Phase I safety trial of intravenous ascorbic acid in patients with severe sepsis. *J Transl Med* 2014;12:32.

29. Dechartres A, Trinquart L, Boutron I, Ravaud P. Influence of trial sample size on treatment effect estimates: meta-epi-

demiological study. *BMJ* 2013;346:f2304.

30. Ferreira-González I, Busse JW, Heels-Ansdell D, et al. Problems with use of composite end points in cardiovascular trials: systematic review of randomised controlled trials. *BMJ* 2007;334:786.

31. Ferreira-González I, Permyer-Miralda G, Busse JW, et al. Methodologic discussions for using and interpreting compos-

ite endpoints are limited, but still identify major concerns. *J Clin Epidemiol* 2007;60:651-7.

32. Fergusson D, Aaron SD, Guyatt G, Hébert P. Post-randomisation exclusions: the intention to treat principle and excluding patients from analysis. *BMJ* 2002;325:652-4.

Copyright © 2022 Massachusetts Medical Society.

RECEIVE IMMEDIATE NOTIFICATION WHEN AN ARTICLE
IS PUBLISHED ONLINE FIRST

To be notified by email when *Journal* articles
are published online first, sign up at NEJM.org.